

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

~~from about 2% to about 90% of a phytochemical concentrate of Commiphora myrrha, Echinesea purpurea and Echinesea augustifolia, said phytochemical concentrate comprising antimicrobial isolates selected from the group consisting of: sesquiterpenes, curzenone, dihydro fuanodien-6-one, 2-methoxyfurandine, elemol, lyndesterene, acetic acid, alpha-amyrone, arabinose, alpha-bisabolene, gamma-bisabolene, cadinene, campesterol, cholesterol, cinnamaldehyde, commiferin, alpha-commiphoric acid, beta-commiphoric acid, gama-commiphoric acid, commiphorinic acid, m-cresol, cumic alcohol, cuminaldehyde, dipentene, elemol, 3-epi-alpha-amyrin, eugenol, furanodiene, furanodienone, galactose, gum, heerabolene, alpha-heerabomyrrhol, beta-heerabomyrrhol, heeraboresene, limonene, 4-O-methyl-glucuronic acid, n-nonacosane, beta-sitosterol, xylose, echinacen, echinacen-B, echinacine, echinacoside, caffeic acid pester, echinolone, enzymes, glucuronic acid, inulin, inuloid, pentadecadiene, polyacetylene compounds, polysaccharides, arabinogalactan, rhamnose, tannins, PSI (a 4-O-methylglucoronarabinoxylan, Mr 35Kd), PSII (an acid rhamnoarbinogalactain, Mr 450 kD), cynarin, 1,5-di-O-caffeoylquinic acid; chloric acid; 2,3-O-di-caffeoyltartaric acid; borneol, borneol acetate; pentadeca-8(z)-en-2-one, germaerene-D; caryophyllene, caryophyllene epoxide; anthocyanin, pyrolizidine alkaloid, lipophilic amide; isobutylamide; polyacetylene; anthocyanin; 3-O-B-D-glucopyranoside; 3-O-(6-O-mabonyl)-B-D-glucopyranoside; tussilagene, isotussilagene, isomeric dodeca-isobutylamide, tetraenoic acid, carophylenes, alkylamides, apigenin, arabinogalacta, ascorbic acid, behenic acid-ethyl acid, betaine, borneol, bornyl acetate, caffeic acid, 2-O-caffeoyl-3-(5-alpha-carboxybeta)-3,4-dihydroxyphenyl, 2-O-caffeoyl-3-O-cumaroyltaric acid, 6-O-caffeoylechinoside, 2-O-caffeoyl-3-O-feruloyltartaric acid, 2-O-caffeoyltartaric acid, calcium, carbonate, beta-carotene, carophyllene, carophyllene epoxide, chloride, chlorogenic acid, chloric acid, chloric acid-methyl ester, cobalt, cyanadin-3-O-(beta-D-glucopyranoside), cynadin-3-(6-O-malonyl-beta-D-glucopyranoside), cynarin, deca-(2e, 4e, 6e)-trienoic acid isobutylamide, des-rhamnosylverbascoside, 3,5-dicaffeoylquinic acid, 4,5-O-dicaffeoylquinic acid, 2,3-O-diferuloyltartaric acid, do-deca-(2e, 4e)-dienoic acid isobutylamide, dodeca-2,4-dien-1-yl isovalerate, dodeca-(2e, 6z, 8e, 10e)-tetraenoic acid isobutylamide, epishobunol, beta-~~

~~farnesene, 2-O-feruloyltartaric acid, germacrene, heptadeca (8z, 11z) dien 2-one, heteroxylan, humulene 8-12, (e)-10-hydroxy-4, 10-dimethyl-4,11-dodecadien 2-one, 13-hydroxyoctadeca (9z, 11e, 15z) trienoic acid, inulin, iron, isochlorogenic acid, isorhamnetin-3-rutinoside, isotussilagine, kaempferol, kaempferol-3-glucoside, kaempferol-3-rutinoside, limonene, luteolin, luteolin-7-glucoside, magnesium, manganese, 2-methyltetradeca-5, 12-diene, 2-methyltetradeca-6, 12-diene, methyl-p-hydroxycinnamate, marcene, niacin, palmitic acid, pentadeca (8z, 11z) dien 2-one, pentadeca (8z, 13z) dien-11-yn 2-one, pentadeca-8-en 2-one, pentadeca (8z)-en 2-one, pentadeca (8z)-en-11, 13-dien 2-one, 1-pentadecene, penta (1, 8z)-diene, phosphorous, alpha-pinene, beta-pinene, polyacetylenes, pontica-epoxide, potassium, protein, quercetagetin-7-glucoside, quercetin, quercetin-3-galactoside, quercetin-3-glucoside, quercetin-3-rutinoside, quercetin-3-xyloside, quercetin-3-xylosylgalactoside, rhamnoarabinogalactan, riboflavin, rutin, rutoside, selenium, silicate, beta-sitosterol, sitosterol-3-beta-D-glucoside, sodium, stigmasterol, sulfate, tartaric acid, tetradeca (8z)-en-11, 13-dien 2-one, thiamin, n-triacontanol, trideca-1-en-3, 5, 7, 9, 10-pentayne, tussilagine, vanillin, verbascoside, and combinations thereof;~~

~~from about 0.005% to about 0.8% quaternary ammonium salt surfactant comprising a member selected from the group consisting of alkyl dimethylbenzylammonium chloride, benzalkonium halide, benzalkonium bromide, benzathonium chloride, alkylbenzyl dimethylammonium chloride, alkyl dimethylbenzylammonium chloride, n-alkyl dimethylbenzylammonium chloride, diisobutylphenoxyethoxyethyl dimethylammonium chloride, n-dimethylbenzylammonium chloride, octyldecyl dimethylammonium chloride, didecyl dimethylammonium chloride, dioctyl dimethylammonium chloride, dialkyl dimethylammonium chloride, octyldecyl dimethylammonium chloride, lauryl dimethylbenzylammonium chloride, o-benzyl-p-chlorophenol, diderly dimethylammonium chloride, doctyl dimethylammonium chloride, alkyl dimethylbenzylammonium chloride, and alkylbenzyl dimethylammonium chloride;~~

sterile water providing a diluent and carrier for said phytochemical concentrate, and the overall ratio of said sterile water to said phytochemical concentrate and said ammonium salt surfactant ranges from about 2:1 to about 100:1; and

from about 0.01% to about 25% of a nutrient comprising folic acid;

from about 40% to about 60% of a phytochemical concentrate of herbaceous botanicals comprising Commiphora myrrha and Echinacea purpurea; said phytochemical concentrate of Commiphora myrrha and Echinacea purpurea providing antimicrobial isolates;

from about 20% to about 60% of an aqueous diluent and carrier for said phytochemical concentrate;

from about 2% to about 12% folic acid providing a nutrient; and

said folic acid cooperating with said Commiphora myrrha and said Echinacea purpurea to treat human immunodeficiency virus;

systemically applying said antimicrobial compound in sufficient concentration in the person infected with human immunodeficiency virus for a sufficient period of time to decrease human immunodeficiency virus in the person;

controlling viral load; and

said antimicrobial isolates of said phytochemical concentrate comprises by weight based upon the total weight of the medical composition:

from about 0.3% to about 9% echinacoside;

from about 0.1% to about 7% PSI (4-O-methylglucoronarabinoxylan, Mr 35 kD) and PSI (acid rhamnoarabinogalactan, Mr 450 kD);

from about 0.1% to about 10% cynarin (1,5-di-o-caffeoylquinic acid) and chloric acid (2,3-O-di-caffeoyltartaric acid) and derivatives thereof;

from about 0.2% to about 4% echinolone;

from about 0.2% to about 8% echinacin B;

from about 0.1 to about 6% echinaceine;

from about 2% to about 7% anthocyanins comprising cyanidin 3-O-B-D-glucopyranoside and 3-O-(6-O-malonyl)-B-D-glucopyranoside;

from about 0.01% to about 0.06% pyrrolizidine alkaloids comprising tussilagine and isotussilagine;

from about 0.003% to about 0.009% isomeric dodeca isobutyalamides and tetroenoic acid; and

Commophora myrrha phytochemicals comprising members selected from the group consisting of: caryophylenes, sesquiterpenes, curzerenone, dihydro fuanodien-6-one; 2-methoxyfuradine, elemol, lyndesterene, acetic acid, alpha-amyrone, arabinose, alpha-bisabolene, gamma-bisabolene, cadinene, campesterol, cholesterol, cinnamaldehyde, commiferin, alpha-commiphoric acid, beta-commiphoric acid, gama-commiphoric acid, commiphorinic acid, m-cresol, cumic alcohol, cuminaldehyde, dipentene, elemol, 3-epi-alpha-amyrin, eugenol, furanodiene, furanodienone, galactose, gum, heerabolene, alpha-heerabomyrrhol, beta-heerabomyrrhol, heeraboresene, limonene, 4-O-methyl-glucuronic acid, n-nonacesane, beta-sitosterol, and xylose.

Claims 32-34 (canceled)

Claim 35 (new): A method for use in treating human immunodeficiency virus, comprising the steps of:

systemically applying an antimicrobial compound providing a medicinal composition into person infected with human immunodeficiency virus;

said antimicrobial compound comprises by weight:

from about 40% to about 60% of a phytochemical concentrate of herbaceous botanicals consisting of Commiphora myrrha and Echinacea purpurea;

from about 20% to about 60% water providing a diluent and carrier for said phytochemical concentrate;

systemically applying said antimicrobial compound into the person infected with human immunodeficiency virus in sufficient concentration and for a sufficient period of time to decrease human immunodeficiency virus in the person;

controlling viral load; and

said antimicrobial isolates of said phytochemical concentrate comprises by weight based upon the total weight of the medical composition:

from about 0.3% to about 9% echinacoside;

from about 0.1% to about 7% PSI (4-O-methylglucoronarabinoxylan, Mr 35 kD) and PSI (acid rhamnoarabinogalactan, Mr 450 kD);

from about 0.1% to about 10% cynarin (1,5-di-o-caffeoylquinic acid) and chlorogenic acid (2,3-O-di-caffeoyltartaric acid) and derivatives thereof;

from about 0.2% to about 4% echinolone;

from about 0.2% to about 8% echinacin B;

from about 0.1 to about 6% echinaceine;

from about 2% to about 7% anthocyanins comprising cyanidin 3-O-B-D-glucopyranoside and 3-O-(6-O-malonyl)-B-D-glucopyranoside;

from about 0.01% to about 0.06% pyrrolizidine alkaloids comprising tussilaginate and isotussilaginate;

from about 0.003% to about 0.009% isomeric dodeca isobutyramides and tetroenoic acid; and

Commiphora myrrha phytochemicals comprising members selected from the group consisting of: caryophyllenes, sesquiterpenes, curzerenone, dihydro farnodien-6-one; 2-methoxyfarnadiene, elemol, lindenene, acetic acid, alpha-amyrone, arabinose, alpha-bisabolene, gamma-bisabolene, cadinene, campesterol, cholesterol, cinnamaldehyde, commiferin, alpha-commiphoric acid, beta-commiphoric acid, gamma-commiphoric acid, commiphorinic acid, m-cresol, cumic alcohol, cuminaldehyde, dipentene, elemol, 3-epi-alpha-amyrin, eugenol, farnodiene, farnodienone, galactose, gum, heerabolene, alpha-heerabomyrrhol, beta-heerabomyrrhol, heerabosene, limonene, 4-O-methyl-glucuronic acid, n-nonacosane, beta-sitosterol, and xylose.

Claim 36. (new) A method for use in treating human immunodeficiency virus in accordance with claim 35, wherein: